

Jiao Tong University School Of Medicine, Shanghai, China, ⁵Hangzhou First People's Hospital, Hangzhou, China

Background: Access to breast cancer screening in China is primarily limited by the shortage of qualified radiologists in primary hospitals. Automated Breast Ultrasound System (ABUS) is a potential method to alleviate current shortages in accessible breast cancer screening. This study aims to evaluate the effectiveness of ABUS by comparing it with Hand Held Ultrasound (HHUS) and Mammography (MAM).

Methods: This study takes place at 3 cancer hospitals and 2 general hospitals. Women aged 30–69, who visited breast surgeons, without visible, suspicious signs of breast cancer and had signed Informed Consent Form were eligible for HHUS and ABUS, and women older than 40 also received MAM. All images were interpreted by qualified doctors based on Breast Imaging Reporting and Data System (BI-RADS). Categories 4–5 were considered to be “suspicious” lesions. The consistency rates and Kappa statistics were calculated to assess the reliability of ABUS compared with HHUS or MAM in each age or hospital group. Participants were divided into four groups by the age of ten, and ABUS was compared with MAM in the older groups (>40 years old) only.

Findings: By taking unilateral breast as the unit of analysis, we have acquired 1734 results for HHUS and ABUS, and 1108 results for MAM. The consistency rates in each age group between HHUS and ABUS were 95.37%, 95.97%, 96.07% and 92.78%, and the Kappa values were 0.79, 0.88, 0.89 and 0.81, respectively; The consistency rates between MAM and ABUS were 92.12%, 92.41% and 93.89%, and the Kappa values were 0.74, 0.77 and 0.83 respectively. In the cancer hospitals, the consistency rates between HHUS and ABUS or MAM were 95.61% and 93.33%, and the Kappa values were 0.87 and 0.81; while in the general hospitals, the consistency rates were 95% and 90.14%, and the Kappa values were lower (0.71 and 0.55).

Interpretation: Reliability was observed when comparisons were made between each age group. ABUS images can be collected by technicians and interpreted by qualified doctors. Considering the absence of qualified radiologists, ABUS may play an important role in general and primary hospitals. Other clinical performance indicators of ABUS, including sensitivity and specificity, need to be further demonstrated.

Source of Funding: Investigator Initiated Research Sponsored By GE Healthcare.

Abstract #: 1.026_WOM

Prevalence of Teenage Pregnancy at Saint-Nicolas Hospital in Saint-Marc, a Community Hospital in Haiti

E.F. Julceus¹, O. Sainterant², R. St-Jean², N. Sobler³, K. Israel²; ¹Zanmi Lasante, Tabarre, Haiti, ²Zanmi Lasante, Port au Prince, Haiti, ³City College of New York, New York, USA

Background: Teenage pregnancy is a significant public health problem. It affects 10% of births worldwide and has harmful health effects on mothers and newborns. Additionally, it is stigmatized in many countries, resulting negative socioeconomic consequences. Several risk factors have been identified, however none of the

previous studies conducted in the Caribbean explored the role of Carnival. Carnival is often thought to be conduit for unplanned sexual encounters. This study aimed to determine the proportion of teen pregnancy at Saint-Nicolas Hospital (HSN) broadly, and at the different points were the pregnant woman (PW) sought care as well as testing whether the proportion increased during the carnival period.

Methods: This cross sectional study included PW seen from October 2013 to September 2014 in HSN for antenatal care, delivery and post abortion care. These women were classified according to the first service received. Teenage pregnancy was defined as pregnancy in a woman under 20 years. Data on their last menstrual period was used to determine the quarter in which the women became pregnant. The carnival period included January to March. Data were extracted from the register, entered into Microsoft excel 2013, and analyzed using Epi Info TM 7 and SPSS 20. We report on proportion of all pregnancies involving teenage pregnancies and used chi-square to test statistical significance of difference in proportions of teenage pregnancies by month of conception and type of service.

Findings: Among 5232 registered pregnancies, 12.82% involved a teen pregnancy. This prevalence was 10.54% among the 2391 woman seen in the delivery service, 14.22% among the 2455 seen for antenatal care, and 18.13% among the 386 seen for post abortion care ($p < 0.0001$). Based on the quarter when women became pregnant, proportion of teen pregnancy varied from 13.34% to 14.84% during the carnival period ($p = 0.97$).

Interpretation: The proportion of teenage pregnancy among all pregnant women seeking services in HSN is slightly higher than the global average. A larger proportion were teenage pregnancies in the post abortion service than the other two, but there was no association with carnival period. Then, it is important to sensitize the population about it throughout the year.

Source of Funding: None.

Abstract #: 1.027_WOM

Rates of Viral Suppression among HIV-positive Women in Rural North-central Nigeria

D. Lee¹, G. Rinombota², U.C. Anaba³, N. Sam-Agudu⁴; ¹University of Maryland, Baltimore, Baltimore, Maryland, USA, ²University of Maryland Baltimore, Baltimore, USA, ³Institution, Federal Capital Territory, Abuja, Nigeria, ⁴University of Maryland Baltimore, Baltimore, MD, USA

Background: The effectiveness of antiretroviral drugs in prevention of mother-to-child transmission (PMTCT) and suppression of viral load (VL) is well-documented. In addition, viral suppression is in line with the 90-90-90 global strategy of HIV elimination. To determine adherence and effectiveness of PMTCT treatment, we assessed VL in postpartum HIV-positive women enrolled in a large PMTCT implementation research study in rural North-Central Nigeria

Methods: Within this prospective cohort study, 497 HIV-positive pregnant women were enrolled from 20 Primary Healthcare Centers (PHCs). Viral load (VL) testing (blood test with lower limit of detection of <20 copies/ml) was performed at 6 months post-partum. Per